



# A range extension of *Gentiana capitata* Buch.-Ham. ex D.Don subsp. *harwanensis* (G.Singh) Halda (Gentianaceae) to Ladakh Trans-Himalaya, India

Mohd Shabir<sup>1,3</sup>, Anzar A. Khuroo<sup>2</sup>, Priyanka Agnihotri<sup>1</sup>, Jay Krishan Tiwari<sup>3</sup>, Tariq Husain<sup>1</sup>

**1** Plant Diversity, Systematics and Herbarium Division, CSIR-National Botanical Research Institute, Lucknow, Uttar Pradesh 226001, India.

**2** Centre for Biodiversity & Taxonomy, Department of Botany, University of Kashmir, Srinagar, Jammu and Kashmir 190006, India. **3** Plant Systematics Laboratory, Department of Botany and Microbiology, HNB Garhwal University, Srinagar Garhwal, Uttarakhand 246174, India.

**Corresponding author:** Mohd Shabir, [shabir1610@gmail.com](mailto:shabir1610@gmail.com).

## Abstract

We collected *Gentiana capitata* subsp. *harwanensis* for the first time from Suru valley in Kargil district of Ladakh region, India, and thus extend this subspecies' geographic distribution to the Trans-Himalayan biogeographic region. This subspecies is endemic to India and so far recorded from the North-west and Western Himalaya of India.

## Key words

Trans-Himalayan biogeographic region; endemic species; new habitat; Suru valley.

**Academic editor:** Amit Kumar | Received 4 September 2018 | Accepted 25 November 2018 | Published 1 February 2019

**Citation:** Shabir M, Khuroo AA, Agnihotri P, Tiwari JK, Husain T (2019) A range extension of *Gentiana capitata* Buch.-Ham. ex D.Don subsp. *harwanensis* (G.Singh) Halda (Gentianaceae) to Ladakh Trans-Himalaya, India. Check List 15 (1): 105–108. <https://doi.org/10.15560/15.1.105>

## Introduction

The genus *Gentiana* L. was first established by Linnaeus in his Species Plantarum (1753). Currently, around the world there are approximately 362 recognized species belonging to 15 sections of the genus (Ho and Liu 2001, Mabberley 2008). In India, *Gentiana* is represented by about 68 species. Of these, approximately 32 species occur in Jammu and Kashmir (Garg 1987, Gupta et al. 2012, Maity 2014, Anilkumar et al. 2015, Shabir et al. 2017, 2018).

*Gentiana capitata* Buch.-Ham. ex D.Don subsp. *harwanensis* Halda was first described by Singh (1976) as *Gentiana harwanensis* G.Singh from Harwan in Kashmir (Jammu and Kashmir) and is endemic to India. The type locality, Harwan (34°09'32" N, 074°54'13" E), is 15 km from Srinagar Kashmir and is within the temperate bio-

geographic region in North-western Himalaya.

Herein, we provide the first record of this subspecies from the Suru valley, in the Kargil district of Ladakh region, India, and extend this subspecies' geographic distribution to the Trans-Himalayan biogeographic region. A detailed description, photographs, and distributional map are included.

## Methods

During a recent botanical trip to Suru valley in Kargil district in June 2018, the first author (MS) came across and collected a specimen of the subspecies reported here. The identification was made by consulting the protologue, the relevant taxonomic literature (Agarwal and Bhattacharayya 1982, Garg 1987, Halda 1995, Ho and Liu 2001, Shabir et al. 2018), and herbarium specimens housed in




**Figure 1.** *Gentiana capitata* Buch.-Ham. ex D.Don subsp. *harwanensis* (G.Singh) Halda. **A.** Habit. **B.** Flower.


the herbaria of Kashmir University (KASH) where the holotype is deposited, National Botanical Research Institute, Lucknow (LWG), and from the image of the isotype deposited in the New York Botanical Garden (NY). The voucher specimens are deposited in LWG.

Morphological characters were observed and pho-

tographs made of the floral parts using a stereo-zoom microscope (Leica). Field photographs were taken by using a camera (Canon EOS D200). Figures were edited using software Adobe Photoshop.



**Figure 2.** *Gentiana capitata* Buch.-Ham. ex D.Don subsp. *harwanensis* (G.Singh) Halda. **A.** Calyx. **B.** Corolla. **C.** Capsule. **D.** Seeds.



**Figure 3.** Map showing the distribution of *Gentiana capitata* Buch.-Ham. ex D. Don subsp. *harwanensis* (G. Singh) Halda in Indian Himalayan region.

## Results

***Gentiana capitata* Buch.-Ham. ex D. Don subsp. *harwanensis* (G. Singh) Halda. In Acta Mus. Richnov. 3: 36 (1995)**

Figures 1, 2

**Type.** Kashmir, Harwan, along drains in forest, 1700 m, 24 April 1969, Gurcharan Singh 1524 (holotype, KASH; isotype, NY- photo!).

*Ciminalis harwanensis* (G. Singh) Omer. in Edinb. J. Bot. 50(1): 67 (1993).

*Gentiana harwanensis* G. Singh, Forest Fl. Srinagar. 126. fig. 5. (1976).

**New record.** India. Ladakh, Trans-Himalayan biogeographic region, Kargil district, Suru valley, riverbed of Suru river at Damsna (34°09'31" N, 075°57'14" E), 3200 m, 10 June 2018, Mohd Shabir, 309945 (LWG) (Fig. 3).

**Other specimens examined.** India. Jammu and Kashmir: along the drains, Harwan, 1700 m, 24 April 1969, Gurcharan Singh 1524 (KASH); same area, 25 July 1970, Gurcharan Singh 1524 (NY). Shallabugh, 1600 m, 02 September 1982, G.H. Dar 4015 (KASH). Along river Suru, Damsna (Kargil) Ladakh, 3200 m, 10 June 2018, Mohd Shabir 309945 (LWG).

**Description.** Annual, 2.5–4.0 cm high. Stem quadrangular, finely lineolate. Basal leaves broadly elliptic, 3.8–5.3 × 2.9–3.7 mm, apex acute, subacuminate, quadriferous,

arranged opposite in 2 series, 3-distinct veins on abaxial surface, mid-vein keeled. Cauline leaves 2–4 pairs, opposite, widely spaced or rarely overlapping, blade 4.4–5.4 × 2.4–3.0 mm, broadly obovate, spathulate, apex acute, recurved, mid-vein keeled abaxially, margin cartilaginous. Inflorescence terminal, clustered, enveloped by upper obovate leaves; pedicel sessile to sub-sessile. Calyx 4.6–5.5 mm long; tube membranous, 3.0–3.5 mm long; lobes 1.5–2.2 × 0.6–0.9 mm, oblong-lanceolate, apex acute, margin membranous, narrowly cartilaginous and crenulate, mid-vein keeled. Corolla 5.4–5.6 mm long; tube 4.5–4.7 mm long; lobes 0.9–1.2 × 0.70–0.85 mm, ovate-oblong, apex acute, margin entire; plicae 0.55–0.75 × 0.45–0.65 mm, ovate, slightly bifid at apex, margin entire. Stamens 5; filaments equal, inserted at the base of corolla tube, 1.34–1.6 mm long; anthers sagittate, ellipsoid, 0.6–0.7 × 0.2–0.3 mm. Style 0.3–0.6 mm long, stigma rounded. Capsule 3.1–5.2 × 1.7–3.5 mm, clavate, obovate and winged distally along the suture. Seeds winged, 0.5–0.6 × 0.20–0.25 mm, brown and surface reticulate.

**Diagnostic characters.** This subspecies is characterized by its stem not naked below, apically branched; basal leaves rosulate, cauline leaves obovate to spathulate, 3 or 4 pairs; flowers sessile to sub-sessile, clustered (3–5) into a terminal head; calyx tube membranous, lobes ovate-oblong, margins narrowly cartilaginous and crenulate; corolla lobes ovate with acute apex; plicae bifid; capsules obovate with their base truncate.



### Key to the infraspecific taxa of *Gentiana capitata*

- 1 Stem naked below, calyx lobes margin purple or white and broadly membranous.....  
.....*G. capitata* subsp. *capitata*
- Stem not naked below, calyx lobes margin narrowly cartilaginous and crenulate.....  
.....*G. capitata* subsp. *harwanensis*

### Discussion

Singh (1976) in his protologue of *Gentiana harwanensis* G.Singh mentioned that the taxon is closely allied to *G. aquatica* L., but differs from the latter in having fasciculate, campanulate flowers into terminal head, almost sessile flowers, obovate to spatulate leaves and bifid plicae. Later, Agarwal et al. (1981) refined the description of *G. harwanensis* because the Singh's (1976) original description did not adequately incorporate the differentiating characters of taxonomic value. Agarwal et al. (1981) also observed that the plant is close to *G. riparia* Kar. & Kir., *G. albicalyx* Burkill and *G. capitata* Buch.-Ham. ex D.Don, but not to *G. aquatica* as observed by Singh (1976). Omer (1993) transferred the taxon to the genus *Ciminalis* Adans., and Halda (1995) gave subspecies rank to *G. harwanensis* (as *G. capitata* subsp. *harwanensis*). Shabir et al. (2017) also treated this taxon as a subspecies.

*Gentiana capitata* subsp. *harwanensis* is endemic to India, where it is known from the submontane and temperate regions in Kashmir, Himachal Pradesh, and Uttarakhand in Indian western Himalaya (Singh 1976, Garg 1987, Gupta et al. 2012, Shabir et al. 2017). The IUCN (2012) categorized it as Data Deficient.

Although major taxonomic explorations have been made in the Ladakh Trans-Himalayan region (Kachroo et al. 1977, Klimeš and Dickore 2005, Chaurasia et al. 2007, Srivastava and Shukla 2015), but there are no reports available on the occurrence of this subspecies in the region. Our report of this subspecies in Suru valley extends its geographic distribution to the Ladakh Himalaya, which is within the Indian Trans-Himalayan biogeographic region.

### Acknowledgements

We thank the Director, CSIR-National Botanical Research Institute, Lucknow, for providing necessary facilities and encouragements. MS also thanks UGC-New Delhi for providing a fellowship to carry out this research work.

### Authors' Contributions

MS collected and photographed the specimens. MS and AAK identified the specimens. MS wrote the manuscript, designed the distribution map and edited the figures. AAK, PA, JKT, and TH analyzed the data, edited and refined the manuscript.

### References

- Agarwal S, Bhattacharyya UC (1982) Studies on *Gentiana capitata* Buch.-Ham. ex D.Don and its allies. Journal of Economic and Taxonomic Botany 3: 981–989.
- Agarwal S, Bhattacharyya UC, Gupta BK (1981) A note on *Gentiana harwanensis* G.Singh. Indian Journal of Forestry 4: 236–238.
- Anilkumar KA, Prabhu Kumar KM, Udayan PS (2015) *Gentiana kurumbae*, a new species of Gentianaceae from the Western Ghats of Kerala, India. Taiwan 60 (2): 81–85. <https://doi.org/10.6165/tai.2015.60.81>
- Chaurasia OP, Ahmad Z, Ballabh B (2007) Ethnobotany and Plants of Trans- Himalaya. Satish Serial Publishing House, Delhi, 544 pp.
- Garg S (1987) Gentianaceae of the North West Himalaya (A Revision). Today & Tomorrow's Printers and Publishers, New Delhi, 342 pp.
- Gupta SA, Mukherjee SK, Mondal M (2012) A census of *Gentiana* L. in India, systematics of flowering plants. In: Maiti G, Mukherjee SK (Eds) Multidisciplinary Approaches in Angiosperm Systematics. University of Kalyani, Kalyani, 53–58.
- Halda JJ (1995) Synopsis of the new system of the genus *Gentiana*. Acta Musei Richnoviensis 3: 3–50.
- Ho TN, Liu S (2001) A Worldwide Monograph of *Gentiana*. Science Press, Beijing, 401 pp.
- IUCN (2012) IUCN Red List Category and Criteria. Version 3.1. IUCN, Gland and Cambridge, 32 pp.
- Kachroo P, Sapru BL, Dhar U (1977) Flora of Ladakh: an Ecological and Taxonomical Appraisal. Bishen Singh and Mahendra Pal Singh, Dehradun, 172 pp.
- Klimeš L, Dickore B (2006) Flora of Ladakh (NW Himalaya)—a Preliminary Check-list. <http://www.butbn.cas.cz/klimes/desert.html>. Accessed on: 2018/11/30.
- Linnaeus C (1753) Species plantarum :exhibentes plantas rite cognitatas, ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas. Tomus 1. Impensis Laurentii Salvii, Holmiae, 227 pp. <https://doi.org/10.5962/bhl.title.669>
- Mabberley DJ (2008) The Plant Book. A Portable Dictionary of Plants, their Classification and Uses, 3rd edition. Cambridge University Press, Cambridge, 354 pp.
- Maity D (2014) A new species of *Gentiana* L. (Gentianaceae) from Sikkim Himalaya. Edinburgh Journal of Botany 71(3): 289–296. <https://doi.org/10.1017/S096042861400016X>
- Omer S, Qaiser M (1993) The genus *Ciminalis* (Gentianaceae) in Pakistan and Kashmir. Edinburgh Journal of Botany 50 (1): 67. <https://doi.org/10.1017/S0960428600000676>
- Shabir M, Agnihotri P, Husain D, Tiwari JK, Husain T (2017) On the current status of the genus *Gentiana* L. (Gentianaceae) in India. Pleione 11 (1): 16–24.
- Shabir M, Agnihotri P, Tiwari JK, Husain T (2018) *Gentiana pringlei* (Gentianaceae) – A new species from cold deserts of Ladakh Himalaya, India. Taiwan 63 (4): 356–359. <https://doi.org/10.6165/tai.2018.63.356>
- Singh G (1976) Forest Flora Srinagar and Plants of Neighbourhood. Bishen Singh and Mahendra Pal Singh, Dehradun, 126 pp.
- Srivastava SK, Shukla AN (2015) Flora of Cold Desert Western Himalaya, India, Vol. 2, Botanical Survey of India, Kolkata, 570 pp.